

# Identification of important habitat for the Pilliga Mouse *Pseudomys pilligaensis*

David C. Paull<sup>1</sup>, David Milledge<sup>2</sup>, Phil Spark<sup>3</sup>, Sally Townley<sup>4</sup> and Kevin Taylor<sup>5</sup>

<sup>1</sup>Office of Environment and Heritage, PO Box 44B, Newcastle, NSW

<sup>2</sup>Landmark Ecological Services, PO Box 100 Suffolk Park, NSW 2481

<sup>3</sup>“Tarcoola”, Back Nundle Rd, Tamworth, NSW 2340

<sup>4</sup>70 Bruxner Park Rd, Korora NSW 2450

<sup>5</sup> 24 Pilot St, Urunga, NSW 2455

Correspondence: [davidcpaull@hotmail.com](mailto:davidcpaull@hotmail.com)

## ABSTRACT

Habitats where rodents breed and congregate provide key refuge and foraging resources and as such may be important for the survival of local populations. The habitat characteristics at spring-time breeding and congregation sites of the Pilliga Mouse *Pseudomys pilligaensis* are analysed here. Data have been gathered from surveys undertaken in 1998-1999 and during a recent survey in 2011. The habitats at the sites (n=11) were found to be shrubby woodland or shrubland dominated by a low shrubby understorey with a projected foliage cover of 20-100%, a groundstorey plant cover of 20-70% and a litter cover of 15-50%. These habitats were found to have floristically diverse understoreys (12-29 species) with a variety of overstorey associations, if present. The results demonstrates that this species occurs in a wider range of vegetation communities than previously reported, though has specific breeding habitat preferences based primarily on understorey and groundstorey conditions. The potential distribution of “important habitat” for *P. pilligaensis* based on these floristic and structural preferences is presented as a predictive map, though it is recognised that the identification of these habitat conditions is best achieved by field examination.

**Key words:** *Pseudomys*, Pilliga, habitat, shrubby, woodland, important

<http://dx.doi.org/10.7882/AZ.2013.010>

## Introduction

*Pseudomys pilligaensis* is a small (10-13 g) murid rodent with a restricted range, currently known only from the Pilliga State Forests, Pilliga East State Conservation Area, Willala Aboriginal Area, Pilliga Nature Reserve and Binnaway Nature Reserve in New South Wales, Australia (NSW Wildlife Atlas, Australian Museum records). The taxonomic status of *P. pilligaensis* is subject to conjecture due to previous work carried out by Ford (2003) though its appearance (rufous pelage, Figure 1), ecological (Paull 2009) and biological aspects of breeding (Prof. B. Breed pers. comm.) indicate unique characteristics. Previous studies undertaken during the last significant La Niña weather event (1998-2001) described this species to be “irruptive” and dispersive (Paull 2006; Tokushima and Jarman 2008; Tokushima *et al.* 2008) with a generalist, but seasonally-shifting diet (Jefferys and Fox 2001; Tokushima and Jarman 2010), and with habitat preferences for low shrubs (30-50% projective foliage cover) and for vegetation associations that include recently burnt gullies, Broombush *Melaleuca uncinata* shrubland, acacia-bloodwood *Acacia burrowii* - *Corymbia trachyphloia* shrubby woodlands and red gum-bloodwood shrubby woodland (Paull 2006; 2009).

Rodents are well documented as having a dynamic population ecology, a strategy which is designed to take advantage of conditions when food is widely available,

particularly after rain (Braithwaite and Brady 1993; Southgate and Masters 1996; Madsen and Shine 1999; Letnic *et al.* 2005). Previous studies of *P. pilligaensis* have shown that animals will use the same sites year round, albeit at very different densities, yet will disappear entirely from some sites (Paull 2006; Tokushima *et al.* 2008; Tokushima



**Figure 1.** A Pilliga Mouse *Pseudomys pilligaensis* trapped during the 2011 survey, showing a rufous tinge to the pelage. Photo: J. McDowell

and Jarman 2008) or appear at sites where they had not been caught previously in the same year (Paull 2006). A similar pattern of low site fidelity has been observed in the closely related *P. novaehollandiae* (Wilson *et al.* 2005) and in *P. fumeus* (Ford *et al.* 2003). As with some other native rodents, autumn densities compared to spring densities can be high in *P. pilligaensis*, with co-habitation, including burrow-sharing, within relatively small areas comprising both sexes and animals of different ages (Paull 2006; Tokushima and Jarman 2008). As well as serving a social function (Ebensperger and Hayes 2008) these high-density autumn sites may be providing an important foraging resource, hypogaeal fungi, a food source which this species relies upon in the cooler months (Jefferys and Fox 2001; Tokushima and Jarman 2010) and which may be spatially limited in distribution.

Tokushima *et al.* (2008) suggest that animals disperse to “refuge sites” in the spring, though spring decline in rodents has also been linked with mortality, a decline in fungal forage and dispersal to breeding territories at this time of year (Cockburn 1981; Rodd and Boonstra 1984). This study focuses on the identification of spring-time breeding and congregation sites and the habitats they support, acknowledging that high-density autumn sites are also important for this species’ year round use of the forest. Together the habitats required to support these essential behavioural patterns and the resources they provide may be regarded as “important” or “critical” habitat. Identification of the potential distribution of important

habitat for *P. pilligaensis* provides key information for the ongoing conservation of this species.

## Methods

Data on the habitat of sites where breeding *P. pilligaensis* have been caught was retrieved from two sources; (1) from unpublished data collected from the central area of the Pilliga East State Forest, New South Wales during surveys conducted in 1998-1999 (Paull 2006) and (2) from data collected during a survey for *P. pilligaensis* conducted in October 2011 on the eastern side of the Newell Highway, Pilliga East State Forest (Milledge 2011). The site locations are depicted in Figure 2. Habitat descriptions of the capture sites (using floristic and structural indices) was undertaken during both survey periods. Structural habitat characteristics of sites were described by identification of the dominant plant species in each stratum together with field estimates of percentage foliage cover at various heights above the ground (<0.5 m, 0.5-2 m, 2-5 m, and overstorey). This was obtained at each site by using a technique where the cover was recorded where it intersected a vertical line over at least 10 points along a trapping transect. Groundstorey cover included grasses, forbs, prostrate plants, mosses, fungi, lichens and cryptogams. Cover of litter and bare ground was also estimated at each point in the same manner. The values for each transect were compiled and averaged. Overall cover preferences for each stratum and groundcover type are expressed as a




Figure 2. Study Area showing two survey locations

mean with a standard deviation and a range.

Trapping methodologies for the 1998-1999 surveys are described in Paull (2009). Surveys were carried out in October (1998) and November (1999). In the 2011 survey, 10 sites were selected to sample a range of vegetation types in the area. At each site, 50 Elliott A-type traps were set for four nights (200 trap nights site<sup>-1</sup>; 2000 trap nights total). Within each transect, trap stations were placed 10 m apart along two parallel transects (with 25 traps per transect) spaced approximately 50 m apart.

During all survey periods, traps were baited with a peanut butter/oats/vanilla essence or truffle oil mixture and set for four consecutive nights. Traps were checked each morning and re-baited when necessary. Captured animals were sexed and were categorised as being adult (>10 g) or sub-adult (<10 g). Animals were marked with a temporary, water soluble die on the rump. Recaptured animals were distinguished from other individuals by comparing their sex and weight. All animals were released at the point of capture.

Vegetation communities where breeding females have been caught were categorised using Benson *et al.* (2010) and then assigned a corresponding Lindsay Forest Type (Lindsay 1967). For mapping purposes, the distribution and extent of these Lindsay Forest Types was estimated using the GIS layer *ForestTypesFNSW\_Plyr*. Some Lindsay Types could only be assigned as areas “to be determined” because they may contain suitable understorey habitat, but there is uncertainty due to a lack of site data in those types. As a measure of correspondence and accuracy with habitat mapping, the NSW Wildlife Atlas locations for *P. pilligaensis* were added to the mapping product.

Breeding habitat is defined as those areas where female *P. pilligaensis* were captured in breeding condition (pregnant, lactating).

Results

In total, 187 individual *P. pilligaensis* were captured between 1998 and 1999 (Paull 2006), including 35

recaptures. A total of five individuals were caught in the spring of 1998 and 12 captured in the spring of 1999, all being breeding females or juvenile animals. Breeding females were detected at seven sites (Table 1). In 2011, 24 individual *P. pilligaensis* were captured (including eight re-captures) and lactating females were caught at three sites (Sites A, B and C).

For all spring survey periods, sites were occupied by single breeding females, except at site Hc in 1999 which was found to contain a lactating and a pregnant female at the same time. At Site K (2011), eight individuals were captured with some recaptures, but without any animals in breeding condition. This site has been included here as the trapping result indicates a degree of site fidelity and spring-time social group of *P. pilligaensis* (Table 1). Sites B (where seven animals were caught) and K were dominated by adult males and juvenile animals. One lactating female was caught at Site B, though was trapped on a different transect to the other animals.

Floristics

Table 2 shows that throughout the three trapping periods, breeding *P. pilligaensis* were captured in different shrubland and shrubby woodland associations. They were assigned vegetation types as per Benson *et al.* (2010). Shrubland/scrub habitats were dominated by *Melaleuca uncinata* (ID 171) with woodland habitat comprising of a mixture of communities including White Bloodwood /Acacia association (*Corymbia trachyphloia* spp. *amphistomatica*/ *Acacia burrowii*) (ID 404); Dirty Gum (Baradine Red Gum *Eucalyptus chloroclada*)-Black Cypress Pine *Callitris endlicheri*-White Bloodwood shrubby woodlands of the Pilliga region (ID 408); Rough-barked Apple *Angophora fibrosa*-red gum-cypress pine woodland on sandy flats mainly in the Pilliga region; (ID 401); and White Bloodwood-Red Ironbark *E. Fibrosa*-cypress pine shrubby sandstone woodlands of the Pilliga region (ID 405). These are depicted in Figures 3-7.

For understorey, the 1998/99 data showed commonly occurring shrub species were *Micromyrtes sessilis*, *Calytrix tetragona*, *Grevillea floribunda*, *Westringia cheelii*, *Eriostemon*

Table 1. Locations of breeding sites for *Pseudomys pilligaensis* (1998-2011)

| Site   | PM | Dem              | Breed | Year | Location (Pilliga East State Forest) | Easting | Northing |
|--------|----|------------------|-------|------|--------------------------------------|---------|----------|
| 2      | 1  | 1 af             | 1     | 1998 | County Line Road                     | 733150  | 6613200  |
| 54     | 4  | 1 af; 2 jf; 1 jm | 1     | 1998 | County Line Road                     | 711100  | 6603350  |
| Aa     | 2  | 1 af             | 1     | 1999 | Reilly's Road                        | 715650  | 6696150  |
| Ca     | 26 | 1 af; 1 jm       | 1     | 1999 | Junction Road                        | 711150  | 6605700  |
| Gb     | 12 | 1 af             | 1     | 1999 | County Line Road                     | 718250  | 6605250  |
| Gc     | 12 | 2 af             | 1     | 1999 | County Line Road                     | 718250  | 6605350  |
| Hc     | 13 | 2 af             | 2     | 1999 | County Line Road                     | 718350  | 6604400  |
| Site A | 1  | 1 af             | 1     | 2011 | X-Line Rd                            | 748237  | 6608034  |
| Site B | 7  | 2 af; 4 am; 1 jf | 1     | 2011 | X-Line Rd                            | 752363  | 6607588  |
| Site C | 4  | 1 af; 3 jf       | 1     | 2011 | Monument Rd                          | 765271  | 6609916  |
| Site K | 8  | 4 am; 2 jm; 2 jf | 0     | 2011 | Brandons Rd                          | 752135  | 6613704  |

PM: Pilliga Mouse individuals caught at site throughout the year (1998/99) and during one spring trapping event (2011); Dem: demographics (a=adult, j=juvenile, f=female, m=male); Breed: Number of breeding females



**Table 2.** Floristic description of vegetation communities where breeding *P. pilligaensis* have been trapped (1998-2011)

| Site           | Overstorey dominants  | Mid-storey dominants  | Understorey dominants   | ID* | SP^   |
|----------------|---|---|---|-----|-------|
| 1998-1999      |   |   |   |     |       |
| Aa<br>Ca       | <i>Melaleuca uncinata</i>   |   | <i>Calytrix tetragona</i><br><i>Micromyrtus sessilis</i><br><i>Grevillea floribunda</i><br><i>Westringia cheelii</i><br><i>Eriostemon ericifolius</i><br><i>Damperia lanceolata</i> | 171 | 18-20 |
| Gb<br>Gc<br>Hc |   |   | <i>Melaleuca uncinata</i><br><i>Micromyrtus sessilis</i><br><i>Grevillea floribunda</i><br><i>Westringia cheelii</i><br><i>Eriostemon ericifolius</i><br><i>Damperia lanceolata</i> | 171 | 15-17 |
| 54             | <i>Corymbia trachyphloia</i> ,<br><i>Acacia burrowii</i>                                  | <i>Dodonaea falcata</i><br><i>Phebalium glandulosum</i>                                 | <i>Hibbertia riparia</i><br><i>Prostanthera leichardtii</i><br><i>Gonocarpus elatus</i><br><i>Actinotus gibbonsii</i>   | 404 | 18    |
| 2              | <i>Corymbia trachyphloia</i> ,<br><i>Eucalyptus chloroclada</i>                           | <i>Callitris endlicheri</i>   | <i>Calytrix tetragona</i> ,<br><i>Grevillea floribunda</i><br><i>Brachyloma daphnoides</i>  | 408 | 19    |
| 2011           |   |   |   |     |       |
| A              | <i>Corymbia trachyphloia</i> ,<br><i>Eucalyptus chloroclada</i>                           | <i>Callitris endlicheri</i><br><i>Cassinia arcuata</i><br><i>Conospermum taxifolium</i> | <i>Calytrix tetragona</i> ,<br><i>Grevillea floribunda</i><br><i>Brachyloma daphnoides</i>  | 408 | 21    |
| B              | <i>Angophora floribunda</i><br><i>Eucalyptus chloroclada</i>                              |   | <i>Boronia glabra</i> ,<br><i>Dodonaea peduncularis</i><br><i>Bossiaea rhombifolia</i>  | 401 | 19    |
| C              | <i>Eucalyptus fibrosa</i><br><i>Corymbia trachyphloia</i><br><i>Eucalyptus chlorocada</i> | <i>Cassinia arcuata</i><br><i>Leptospermum parviflorum</i>                              | <i>Brachyloma daphnoides</i> ,<br><i>Hibbertia obtusifolia</i><br><i>Cymbopogon</i> sp.   | 405 | 29    |
| K              | <i>Eucalyptus chloroclada</i>   | <i>Philotheca salsolifolia</i>  | <i>Aotus mollis</i> ,<br><i>Boronia glabra</i><br><i>Brachyloma daphnoides</i>  | 408 | 12    |

\*after Benson et al. (2010); ^understorey plant species richness



**Figure 3.** Low woodland dominated by White Bloodwood *Corymbia trachyphloia* and Kurricabah *Acacia burrowii* with a dense lower-storey containing a variety of woody shrubs, where breeding *P. pilligaensis* were captured in 1998 (Benson ID 404). Photo: by D. Paull



**Figure 4.** Dense post-fire regrowth shrubland, dominated by Broombush *Melaleuca uncinata*, with *Micromyrtus sessilis*, *Grevillea floribunda*, *Westringia cheelii*, *Eriostemon ericifolius* and *Damperia lanceolata* where breeding and over-wintering *P. pilligaensis* were captured in 1999 (Benson ID 171). Photo: D. Paull





**Figure 5.** Site A where a lactating *Pseudomys pilligaensis* was captured, showing *Corymbia trachyphloia*-*Eucalyptus chloroclada* dominated woodland with a dense lower-storey dominated by *Calytrix tetragona*, *Grevillea floribunda* and *Brachyloma daphnoides* (Benson ID 408). Photo: A. McKinley



**Figure 6.** Site B where seven *P. pilligaensis* were captured including a lactating female and sub-adult animals, showing *Angophora floribunda*-*Eucalyptus chloroclada* dominated woodland with a dense lower-storey dominated by *Boronia glabra*, *Dodonaea peduncularis* and *Bossiaea rhombifolia* (Benson ID 401). Photo: A. McKinley



**Figure 7.** Site C where four *P. pilligaensis* were captured including a lactating female and sub-adults, showing *Eucalyptus fibrosa*-*Corymbia trachyphloia*-*E. chloroclada* dominated woodland with a dense lower-storey dominated by *Schoenus ericetorum*, *Hibbertia obtusifolia* and *Cymbopogon* sp. (Benson ID 405). Photo: A. McKinley

*ericifolius* and *Damperia lanceolata*. In 2011, dominant understorey species were most commonly *Boronia glabra*, *Brachyloma daphnoides*, *C. tetragona* and *G. floribunda*. The latter two were found to be dominant at sites in both 1998/99 and in 2011. Sites where *P. pilligaensis* was recorded were found to have an understorey plant diversity of between 15 to 20 plant species in 1998/99 and 12-29 species in 2011. Sites burnt in the last two years prior to trapping (Gb, Gc, Hb and K) tended to have a lower plant diversity (Table 2).

### Vegetation Structure

Table 3 shows the structural characteristics of all sites considered together ( $n=11$ ). They were found to have a ground combine storey plant cover of  $35.9\% \pm 14.9$  (range = 20-70%) and an understorey shrub cover of  $55\% \pm 22.7$  (range = 20-100%). These sites had a bare ground cover of  $30\% \pm 19.5$  (range = 0-60 %), with a litter cover of  $34.1\% \pm 10.6$  (range = 20-50%). Both overstorey (range 0-60%) and midstorey (0-70%) are not useful habitat predictors due to their high level of variance compared to the mean cover and range values (Table 3). Site data suggests no difference in understorey cover between burnt and unburnt sites, though unburnt sites tend to have less bare ground and a higher level of litter cover.

### Mapping the distribution of important habitat

To facilitate mapping of the potential distribution of important habitat for this species, vegetation communities where breeding animals have been caught were placed within the grouped Lindsay Types (Lindsay 1967) of "Apple/Red Gum", "Broad leaf Ironbark/Bloodwood", "Red Gum/Broad leaf Ironbark/Bloodwood", "Broombush" and "Acacia/Bloodwood" (Table 4). Using the digitised Lindsay Type Forests NSW mapping layer, the extent of "potential habitat" (106,839 ha) and habitat "to be determined" (26,603 ha) across the Pilliga East State Forest and Pilliga East State Conservation Area is indicated in Figure 8. The Lindsay Forest Types "Narrow leaf Ironbark-Bloodwood-Red Gum" associations (23,826 ha) and others described as "Heavy Undergrowth" or "Scrub" (2,777 ha) may contain important habitat for this species but have not been investigated.

## Discussion

### Population density and phenology

In 2011, the average spring-time density of individuals encountered was  $3.57 \pm 3.05$  individuals/site ( $n=24$ , range 1-8) while in 1998/99 the average density was found to be lower at  $1.88 \pm 0.99$  individuals/site ( $n=13$ , range=1-4). This is similar to overall spring density found by Tokushima *et al.* (2008) who recorded densities of 0-2 individuals  $\cdot \text{ha}^{-1}$ , though one site (2A) recorded 9 individuals/site in October 2000, comprising of mostly males. Both studies show a spring-time social organisation comprising of solitary breeding females and social groups of males and non-reproductive animals at lower densities than in the autumn.

Studies reported hererecorded breeding in October and November which is consistent with the known breeding season of this species, though this time of year is also a time of dispersal (Paull 2006; Tokushima and Jarman 2008).

**Table 3.** Habitat structure at sites where breeding *P. pilligaensis* were trapped (1998-2011)

| site | UH | UC    | MH | MC    | LH  | LC    | BG    | LL    | GC    | Burnt |
|------|----|-------|----|-------|-----|-------|-------|-------|-------|-------|
| Aa   | 0  | -     | 2  | 40    | <1  | 50    | 10    | 40    | 50    | No    |
| Ca   | 0  | -     | 2  | 70    | <1  | 20    | 0     | 30    | 70    | No    |
| Gb   | 0  | -     | -  | -     | <1  | 70    | 20    | 40    | 40    | Yes   |
| Gc   | 0  | -     | -  | -     | <1  | 100   | 50    | 30    | 20    | Yes   |
| Hc   | 0  | -     | -  | -     | <1  | 70    | 60    | 20    | 20    | Yes   |
| 2    | 10 | 20    | 2  | 20    | <1  | 70    | 30    | 40    | 30    | No    |
| 54   | 4  | 60    | 2  | 10    | <1  | 60    | 0     | 50    | 50    | No    |
| A    | 15 | 20    | 3  | 20    | <1  | 50    | 40    | 30    | 30    | No    |
| B    | 15 | 20    | 0  | 0     | <1  | 60    | 40    | 30    | 30    | No    |
| C    | 12 | 15    | 2  | 50    | 0.5 | 20    | 30    | 50    | 20    | No    |
| K    | 12 | 15    | 2  | 50    | 0.5 | 35    | 50    | 15    | 35    | Yes   |
| Mean |    | 13.64 |    | 23.64 |     | 55    | 30    | 34.09 | 35.91 |       |
| SD   |    | 17.07 |    | 23.85 |     | 22.66 | 19.54 | 10.62 | 14.9  |       |

**Key:** UH: Upper vegetation (canopy) height (m); UC: Upper vegetation % foliage cover; MH: Mid-storey vegetation height (m); MC: Mid vegetation % foliage cover; LH: Under-storey Vegetation height (m); LC: Lower Vegetation % foliage cover; BG: Bare ground % cover; LL: Leaf litter % cover; GC: Ground plant % cover.

**Table 4.** Extent of known breeding habitat associations for *P. pilligaensis* within the Pilliga forests

| Overstorey association                          | Low shrub cover | Ground plant cover | Lindsay Type                            | Pilliga Extent (ha)* | Source                      |
|---|-----------------|--------------------|---|----------------------|-----------------------------|
| <i>A. floribunda</i> , red gum                  | 60 %            | 30%                | Rough bark Apple/ Red Gum               | 30,879.5             | Milledge 2011               |
| <i>E. fibrosa</i> , <i>C. trachyphloia</i>      | 20-50 %         | 20-30%             | Broad leaf Ironbark/ Bloodwood          | 45,902.6             | Milledge 2011               |
| <i>C. trachyphloia</i> , <i>E. chloroclada</i>  | 30-50 %         | 30-35%             | Red Gum/ Broad leaf Ironbark/ Bloodwood | 5,128.6              | Milledge 2011<br>Paull 2006 |
| <i>Melaleuca uncinata</i> ,                     | 20-100 %        | 20-70%             | Broombush                               | 23,896.6             | Paull 2006                  |
| <i>C. trachyphloia</i> , <i>Acacia burrowii</i> | 60 %            | 50%                | Acacia/ Bloodwood                       | 1,030.6              | Paull 2006                  |
| <b>Total</b>                                    |                 |                    |   | <b>106 837.9</b>     |                             |

\* Lindsay (1967)

Capture of non-reproductive adult females at the same time as reproductive females at different sites confirms that breeding is not entirely synchronous in *P. pilligaensis*, with a breeding season known to extend from October to April (Fox and Briscoe 1980; Paull 2006; Tokushima et al. 2008).

### Breeding ecology

While the 1998/99 surveys showed that breeding habitat for this species was mostly shrublands/scrubs, the 2011 surveys confirmed that breeding habitat for this species includes woodland vegetation communities with overstories which include *A. floribunda*, *E. chloroclada* and *E. fibrosa* with a moderate to high understorey cover. Studies undertaken by Tokushima et al. (2008) found high numbers of *P. pilligaensis* at four sites, with breeding also recorded at some of these sites, but did not record details of their floristics or structure. They did note in general a “sparse woodland dominated by *Eucalyptus*, *Callitris* and *Angophora* species, with a diverse mid- and understorey of woody shrubs, and a very sparse vegetation, although often much litter, covering the ground”.


What may be considered as being a “sparse”, or a “moderate”, density is unclear, however, the results presented here show that foliage cover at 0.5-1 m in height varied considerably between sites from a relatively low cover (20%) to a high cover (100%). What is consistent in the data collected from 1998/99 and 2011 is the dominance of woody shrubs in the understorey, the high shrub diversity and in some instances the species composition of the understorey.

Despite the two study areas being a considerable distance apart (~50 km). Some common understorey dominants in the two study areas were the myrtaceous shrub *Calytnix tetragona*, the proteacid *Grevillea floribunda* and the Epacrid heath *Brachyloma daphnoides*. Tokushima et al. (2008) recorded common understorey species of *Boronia*, *Gahnia*, *Daviesia*, *Pomax* and *Cassinia* at their study sites in the Pilliga Nature Reserve.

While sites dominated by Broombush *M. uncinata* sustained breeding females and all-year occupancy in 1999, others were trapped without success in both survey periods. This inconsistency may in part be explained by the fire histories of Broombush habitats. Paull (2009) found that only young (1.5- 3 yr old regrowth) and mature (>25 years old) Broombush sites were selected preferentially by *P. pilligaensis*. The age of the Broombush vegetation in the 2011 study was six years since the last wildfire and displayed a growth form with a dense projective foliage cover at 2 metres in height conditions which *P. pilligaensis* was found to significantly avoid.

When compared to species to which it is most closely related, *P. pilligaensis* selects more restricted habitats, perhaps due to a more restricted geographical distribution. *P. delicatulus* has been recorded as breeding in tropical savannahs (Braithwaite and Brady 1993) and coastal sand-dunes (Taylor and Horner 1970). Taylor and Horner (1970) state *P. delicatulus* avoids monsoon forest or woodlands, however other studies have caught this





**Figure 8.** Distribution of important habitat of *Pseudomys pilligaensis* showing predicted areas (green) and areas to be determined (yellow). NSW Wildlife Atlas records of *P. pilligaensis* are shown in red, and records from 2011 survey are shown in black.

species in tropical *Eucalyptus* woodlands, though breeding is not identified (Kerle and Burgmann 1984; Woinarski *et al.* 2001). *P. novaehollandiae* is known to breed in coastal grasslands, heathlands, regenerating pine plantations and shrubby woodlands (Kemper 1980; 1990; Fox and Fox 1978; 1984; Lock and Wilson 1999), including vegetated coastal dune habitats (Wilson *et al.* 2005). In terms of its habitat selection, *P. pilligaensis* is most like other temperate zone *Pseudomys* species such as *P. novaehollandiae* and *P. fumeus* which also show a preference for habitats with understories dominated by woody shrubs.

### Distribution of important habitat

Records of this species obtained from the NSW Wildlife Atlas have been added as a measure of correspondence with the predictive habitat mapping in Figure 8 and despite some outliers to the west, the mapping indicates a reasonably close match between the distribution of records and predicted distribution of important habitat. However it should be noted that not all records represent congregations or breeding sites and some individual records may be represent dispersing animals.

While the extent of possible habitat covers some than 106,800 ha in the Pilliga State Forest and State Conservation Areas at any one time, not all of this area would actually be suitable as important habitat due to local floristic variation and different disturbance and fire histories across the forest. A preliminary field investigation of habitat suitability (Milledge 2011) showed that within a broad area identified as possible habitat only 20% on average was suitable for *P. pilligaensis*. As this species, like other *Pseudomys*, is primarily dependent upon under and ground storey conditions which are regulated by seasonal, stochastic and climatic factors, mapping of important habitat cannot be undertaken accurately without a field assessment of the existing understory conditions using the criteria identified here.

The identification and extent of important habitat for this species is essential for conducting ongoing monitoring of *P. pilligaensis* in order to track longer term population trends, particularly in the context of drought, climate change and coal seam gas production and warrants further investigation in order that predictive modelling can be refined.

### Acknowledgements

Many people too numerous to name contributed to the success of this study, particularly members of the Northern Inland Council for the Environment, Coonabarabran and Upper Castlereagh Catchment and Landcare Group and The Wilderness Society in providing field support. We especially wish to thank Carmel Flint for organisation, Darren Bailey, Annette McKinley, Hugh and Nan Nicholson and Kate Smilie

for site flora assessments and Arnold Dobbe, Steve Rotschild, Glenn Michelle, Dale Bridge, Georgia Beyer and Justin McDowell for assistance with setting and checking traps.

This study was conducted under the Forests NSW Special Permit for Research XX51747; NSW Scientific Licence S12398 and Animal Care and Ethics approval; File No. 07/4017.

## References

- Benson, J.S., Richards, P., Waller, S. & Allen, C. 2010. New South Wales Vegetation Classification and Assessment: Part 3 Plant communities in the NSW North Western Slopes and west-New England region and update of Western Plains and NSW South Western Slopes Bioregion. Version 3 of the NSWVCA database. *Cunninghamia* 11(4): 457-579.
- Braithwaite, R.W. and Brady, P. 1993. The delicate mouse, *Pseudomys delicatulus*: a continuous breeder waiting for the good times. *Australian Mammalogy* 16: 93-96.
- Cockburn, A. 1981. Population regulation and dispersion of the smoky mouse, *Pseudomys fumeus* I. Dietary determinants of microhabitat preference. *Australian Journal of Ecology* 6(3): 231-254. <http://dx.doi.org/10.1111/j.1442-9993.1981.tb01574.x>
- Ebensperger, L.A. and Hayes, L.D. 2008. On the dynamics of rodent social groups. *Behavioural Processes* 79 (2): 85-92. <http://dx.doi.org/10.1016/j.beproc.2008.05.006>
- Ford, F. (2003). Conilurine rodent evolution: the role of ecology in modifying the consequences of evolutionary change. PhD Thesis, Australian National University.
- Ford, F., Cockburn, A. and Broome, L. 2003. Habitat preference, diet and demography of the smoky mouse *Pseudomys fumeus* (Rodentia: Muridae) in south-eastern New South Wales. *Wildlife Research* 30: 89-101. <http://dx.doi.org/10.1071/WR01092>
- Fox, B.J. and Fox, M.D. 1978. Recolonization of coastal heath by *Pseudomys novaehollandiae* (Muridae) following sand mining. *Australian Journal of Ecology* 3: 447-65. <http://dx.doi.org/10.1111/j.1442-9993.1978.tb01191.x>
- Fox, B.J. and Fox, M.D. 1984. Small mammal recolonisation of open forest following sand mining. *Australian Journal of Ecology* 9: 241-52. <http://dx.doi.org/10.1111/j.1442-9993.1984.tb01361.x>
- Fox, B. J. and Briscoe, D. A., 1980. *Pseudomys pilligaensis*, a new species of murid rodent from the Pilliga Scrub, northern New South Wales. *Aust. Mammal.* 3: 109-26.
- Jefferys, E.A. & Fox, B.J. 2001. The diet of the Pilliga Mouse, *Pseudomys pilligaensis* (Rodentia: Muridae) from the Pilliga Scrub, northern New South Wales. *Proceedings of the Linnean Society of New South Wales* 123: 89-99.
- Kemper, C.M. 1980. Reproduction of *Pseudomys novaehollandiae* (Muridae) in the wild. *Australian Wildlife Research* 7: 385-4012. <http://dx.doi.org/10.1071/WR9800385>
- Kemper, C.M. 1990. Small mammals and habitat disturbance in open forest of coastal New South Wales. I. Population parameters. *Australian Wildlife Research* 17: 195-206. <http://dx.doi.org/10.1071/WR9900195>
- Kerle, J.A. and Burgmann, M.A. 1984. Some aspects of the ecology of the mammal fauna of the Jabiluka Area, Northern Territory. *Australian Wildlife Research* 11: 207-22. <http://dx.doi.org/10.1071/WR9840207>
- Letnic, M., Tamayo, B., and Dickman, C. R. 2005. The responses of mammals to La Niña (El Niño southern oscillation)-associated rainfall, predation, and wildfire in central Australia. *Journal of Mammalogy* 86: 689-703. [http://dx.doi.org/10.1644/1545-1542\(2005\)086\[0689:TROMTL\]2.0.CO;2](http://dx.doi.org/10.1644/1545-1542(2005)086[0689:TROMTL]2.0.CO;2)
- Lock, M.L. and Wilson, B.A. 1999. The distribution of the New Holland Mouse (*Pseudomys novaehollandiae*) with respect to vegetation near Anglesea, Victoria. *Wildlife Research* 26: 565-577. <http://dx.doi.org/10.1071/WR97050>
- Lindsay, A.D. 1967. Forest types of the New South Wales cypress pine zone. Technical Paper No. 8. Forestry Commission of NSW; Sydney.
- Madsen, T. and Shine, R. 1999. Rainfall and Rats: Climatically-driven dynamics of a tropic rodent population. *Australian Journal of Ecology* 24: 80-89. <http://dx.doi.org/10.1046/j.1442-9993.1999.00948.x>
- Milledge, D.R. 2011. *National Significance: The Ecological Values of Pilliga East Forest and the Threats Posed by Coal Seam Gas Mining 2011-2012*. A report prepared for the Northern Inland Council for the Environment and the Coonabarabran and Upper Castlereagh Catchment and Landcare Group.
- Paull, D.C. 2006. The biology and ecophysiology of the Pilliga Mouse *Pseudomys pilligaensis* and the Common Dunnart *Sminthopsis murina* in Pilliga East State Forest. Master of Science Thesis – Zoology, University of New England, Armidale, NSW.
- Paull, D.C. 2009. Habitat and post-fire selection of the Pilliga Mouse *Pseudomys pilligaensis* in Pilliga East State Forest. *Pacific Conservation Biology* 15: 254-267.
- Rodd, F.H. and Boonstra, R. 1984. The spring decline in the meadow vole, *Microtus pennsylvanicus*: the effect of density. *Canadian Journal of Zoology*, 1984, 62(8): 1464-1473. <http://dx.doi.org/10.1139/z84-212>
- Southgate, R. and Masters, P. 1996. Fluctuations of rodent populations in response to rainfall and fire in a central Australian hummock grassland dominated by *Plectrachne schinzii*. *Wildlife Research* 23, 289-303. <http://dx.doi.org/10.1071/WR9960289>
- Taylor, J.M. and Horner, B.E. 1970. Observations on Reproduction in Leggadina (Rodentia, Muridae). *Journal of Mammalogy* 51(1): 10-17. <http://dx.doi.org/10.2307/1378526>
- Tokushima, H., Green, S.W. and Jarman, P. 2008. Ecology of the rare but irruptive Pilliga mouse (*Pseudomys pilligaensis*). I. Population fluctuation and breeding season. *Australian Journal of Zoology* 56: 363-373. <http://dx.doi.org/10.1071/ZO08042>
- Tokushima, H. & Jarman, P. 2008. Ecology of the rare but irruptive Pilliga mouse (*Pseudomys pilligaensis*). II. Demography, home range and dispersal. *Australian Journal of Zoology* 56: 375-387. <http://dx.doi.org/10.1071/ZO08043>
- Tokushima, H. & Jarman P. 2010. Ecology of the rare but irruptive Pilliga mouse, *Pseudomys pilligaensis*. III. Dietary ecology. *Australian Journal of Zoology*, 58: 85-93. <http://dx.doi.org/10.1071/ZO09107>
- Wilson, B. A., White, N. M., Hanley, A., and Tidey, D. L. 2005. Population fluctuations of the New Holland mouse *Pseudomys novaehollandiae* at Wilson's Promontory National Park, Victoria. *Australian Mammalogy* 27: 49-60. <http://dx.doi.org/10.1071/AM05049>
- Woinarski, J.C.Z., Milne, D.J. and Wangeneen, G. 2001. Changes in mammal populations in relatively intact landscapes of Kakadu National Park, Northern Territory, Australia. *Austral Ecology* 26: 360-370. <http://dx.doi.org/10.1046/j.1442-9993.2001.01121.x>